

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

is illustrated by many very appropriate photographs. The form and structure of trees are also carefully considered. The second part is devoted to a manual of the trees of the state, and is well equipped with keys, glossary, and illustrative drawings. A noticeable feature of the illustrations of the individual species is the drawing of the buds on a large scale. It is safe to say that it will take a first rank among the numerous tree manuals now available.—Geo. D. Fuller.

Montane plants of the Rocky Mountains.—RYDBERG,²³ in continuing his studies of the flora of the Rocky Mountains, has added to the articles already noted in this journal²⁴ an investigation of the distribution of the montane species. He finds about 1900 species in this zone, of which one-half are to be regarded as typical inhabitants of this area. Less than 15 per cent are transcontinental, while 53 per cent are endemic. A close analysis is made of the constituents of the flora peculiar to the northern and southern portions of the region as contrasted with that common to both.—Geo. D. Fuller.

Sedge associations in Colorado.—In studying the sedges of northern Colorado, Ramaley25 shows that the genus Carex not only is of decided importance, but that species of this genus dominate many plant associations, particularly in the montane, subalpine, and alpine regions. These associations are either hydrophytic or xerophytic in character, and represent early stages in succession, for as mesophytism is approached the sedges are replaced by grasses and dicotyledons. The principal associations involved are briefly described and their sedge components noted. Of the 44 species of Carex listed, 20 are classed as hydrophytic, 15 as xerophytic, and 9 only as mesophytic.—Geo. D. Fuller.

New African plants.—Engler,²⁶ in continuation of his studies of the African flora, has described 45 new species of Sterculiaceae, 40 of which belong to *Hermannia*, 29 new species of Guttiferae, and 3 new species of Violaceae (belonging to *Hybanthus*).—J. M. C.

A new genus of Umbelliferae.—Thellung²⁷ has described a new genus (Scandicium) of Umbelliferae from the Mediterranean steppe region and Western Asia, based on Scandix stellata Solander. In addition to the species, numerous varieties are described.—J. M. C.

²³ RYDBERG, P. A., Phytogeographical notes on the Rocky Mountain region. VIII. Distribution of the montane plants. Bull. Torr. Bot. Club 46:295-327. 1919.

²⁴ Bot. Gaz. **62**:83-84. 1916; **63**:423-424. 1917; **65**:195. 1918.

²⁵ RAMALEY, FRANCIS, The rôle of sedges in some Colorado plant communities. Amer. Jour. Bot. 6:120–130. fig. 2. 1919.

²⁶ ENGLER, A., Beiträge zur Flora von Afrika. XLVII. Bot. Jahrb. **55**:350–400. **1919**.

²⁷ THELLUNG, A., *Scandicium*, ein neues Umbelliferen-Genus. Sonderabdruck aus Fedde, Repertorium 16:15-22. 1919.